

# Probability Theory

CMPUT 366: Intelligent Systems

P&M §8.1

# Logistics & Assignment #1

- **Midterm** is **March 9** (see eClass for other important dates)
- **Assignment #1 was released on Monday**  
See eClass
- Due **February 3** ~~before lecture~~ **at 11:59pm**
- This week's lab: **Thursday, 5:00pm to 8:00pm, BS M 149**  
(this room!)
  - Not mandatory
  - You can get help from the TAs on your assignment in labs
  - Brief Python refresher at the beginning

# Recap: Search

- Agent searches **internal representation** to find solution
- **Fully-observable, deterministic, offline, single-agent** problems
- **Graph search** finds a **sequence of actions** to a goal node
  - Efficiency gains from using **heuristic functions** to encode **domain knowledge**
- **Local search** finds a goal node by repeatedly making **small changes** to the current state
  - **Random steps** and **random restarts** help handle **local optima, completeness**

# Lecture Outline

1. Recap
2. Uncertainty
3. Probability Semantics
4. Conditional Probability
5. Expected Value

# Uncertainty

- In search problems, agent has **perfect knowledge** of the world and its dynamics
- In most applications, an agent cannot just **make assumptions** and then act according to those assumptions
- Knowledge is **uncertain**:
  - Must consider **multiple** hypotheses
  - Must **update** beliefs about which hypotheses are likely given **observations**

# Example: Wearing a Seatbelt

- An agent has to decide between **three actions**:
  1. Drive without wearing a seatbelt
  2. Drive while wearing a seatbelt
  3. Stay home
- If the agent thinks that an accident **will** happen, it will just stay home
- If the agent thinks that an accident **will not** happen, it will not bother to wear a seatbelt!
- Wearing a seatbelt only makes sense because the agent is **uncertain** about whether driving will lead to an accident.

# Measuring Uncertainty

- **Probability** is a way of **measuring** uncertainty
- We assign a number between 0 and 1 to **events** (hypotheses):
  - **0** means absolutely certain that statement is **false**
  - **1** means absolutely certain that statement is **true**
  - **Intermediate** values mean more or less certain
- Probability is a measurement of **uncertainty**, **not truth**
  - A statement with probability .75 is not "mostly true"
  - Rather, we **believe** it is more **likely** to be true than not

# Subjective vs. Objective: The Frequentist Perspective

- Probabilities can be interpreted as **objective** statements about the **world**, or as **subjective** statements about an agent's **beliefs**.
- Objective view is called **frequentist**:
  - The probability of an event is the proportion of times it would happen **in the long run** of **repeated experiments**
  - Every event has a single, **true** probability
  - Events that can only happen **once** don't have a well-defined probability



# Subjective vs. Objective: The Bayesian Perspective

- Probabilities can be interpreted as **objective** statements about the **world**, or as **subjective** statements about an agent's **beliefs**.
- Subjective view is called **Bayesian**:
  - The probability of an event is a measure of an agent's **belief** about its likelihood
  - Different agents can legitimately have **different beliefs**, so they can legitimately assign **different probabilities** to the same event
  - There is **only one way** to **update** those beliefs in response to new data
- In this course, we will primarily take the **Bayesian** view

# Example: Dice

- Diane rolls a **fair, six-sided die**, and gets the number  $X$ 
  - **Question:** What is  $P(X = 5)$ ? (the probability that Diane rolled a 5)
- Diane truthfully tells Oliver that she rolled an **odd** number.
  - **Question:** What should **Oliver** believe  $P(X = 5)$  is?
- Diane truthfully tells Greta that she rolled a number  $\geq 5$ .
  - **Question:** What should **Greta** believe  $P(X = 5)$  is?
- **Question:** What is  $P(X = 5)$ ?

# Semantics: Possible Worlds

- **Random variables** take values from a **domain**.  
We will write them as uppercase letters (e.g.,  $X, Y, D$ , etc.)
- A **possible world** is a **complete assignment** of values to variables  
We will usually write a single "world" as  $\omega$  and the set of all possible worlds as  $\Omega$
- A **probability measure** is a function  $P : \Omega \rightarrow \mathbb{R}$  over **possible worlds**  $\omega$  satisfying:

1. 
$$\sum_{\omega \in \Omega} P(\omega) = 1$$

2. 
$$P(\omega) \geq 0 \quad \forall \omega \in \Omega$$

# Propositions

- A **primitive proposition** is an equality or inequality expression  
E.g.,  $X = 5$  or  $X \geq 4$
- A **proposition** is built up from other propositions using **logical connectives**.  
E.g.,  $(X = 1 \vee X = 3 \vee X = 5)$
- The **probability** of a proposition is the sum of the probabilities of the possible worlds in which that proposition is true:

$$P(\alpha) = \sum_{\omega: \omega \models \alpha} P(\omega) \quad \omega \models \alpha \text{ means "}\alpha \text{ is true in } \omega\text{"}$$

- Therefore:

$$P(\alpha \vee \beta) \geq P(\alpha)$$

$\alpha \vee \beta$  means " $\alpha$  OR  $\beta$ "

$$P(\alpha \wedge \beta) \leq P(\alpha)$$

$\alpha \wedge \beta$  means " $\alpha$  AND  $\beta$ "

$$P(\neg \alpha) = 1 - P(\alpha)$$

$\neg \alpha$  means "NOT  $\alpha$ "

# Joint Distributions

- In our dice example, there was a **single** random variable
- We typically want to think about the interactions of **multiple** random variables
- A **joint distribution** assigns a probability to each full assignment of values to variables
  - e.g.,  $P(X = 1, Y = 5)$ . Equivalent to  $P(X = 1 \wedge Y = 5)$
  - Can view this as another way of specifying a single **possible world**

# Joint Distribution Example

- What might a day be like in Edmonton?  
Random variables:
  - **Weather**,  
with domain {clear, snowing}
  - **Temperature**,  
with domain {mild, cold, very\_cold}
- **Joint distribution**  
 $P(\text{Weather}, \text{Temperature})$ :

Weather	Temperature	P
clear	mild	0.20
clear	cold	0.30
clear	very cold	0.25
snowing	mild	0.05
snowing	cold	0.10
snowing	very cold	0.10

# Marginalization

## Question:

What is the **marginal distribution** of Weather?

- **Marginalization** is using a joint distribution  $P(X_1, \dots, X_m, \dots, X_n)$  to compute a distribution over a smaller number of variables  $P(X_1, \dots, X_m)$ 
  - Smaller distribution is called the **marginal distribution** of its variables
- We compute the marginal distribution by summing out the other variables:

$$P(X, Y) = \sum_{z \in \text{dom}(Z)} P(X, Y, Z = z)$$

Weather	Temperature	P
clear	mild	0.20
clear	cold	0.30
clear	very cold	0.25
snowing	mild	0.05
snowing	cold	0.10
snowing	very cold	0.10

# Conditional Probability

- Agents need to be able to **update** their beliefs based on new **observations**
- This process is called **conditioning**
- We write  $P(h \mid e)$  to denote "probability of **hypothesis**  $h$  given that we have observed **evidence**  $e$ "
  - $P(h \mid e)$  is the **probability of  $h$  conditional on  $e$**



# Semantics of Conditional Probability

- Evidence  $e$  lets us **rule out** all of the worlds that are incompatible with  $e$ 
  - E.g., if I observe that the weather is clear, I should no longer assign **any** probability to the worlds in which it is snowing
- We need to **normalize** the probabilities of the remaining worlds to ensure that the probabilities of possible worlds sum to 1

$$P(\omega | e) = \begin{cases} \frac{1}{P(e)} P(\omega) & \text{if } \omega \in e, \\ 0 & \text{otherwise.} \end{cases}$$

# Conditional Probability Example

- My initial marginal belief about the weather was:  
 $P(\textit{Weather} = \textit{snow}) = 0.25$
  - Suppose I observe that the temperature is **mild**.
  - **Question:** What should I **now** believe about the weather?
1. **Rule out** incompatible worlds
  2. **Normalize** remaining probabilities

Weather	P
clear	$.20 / (.20 + .05) = 0.8$
snowing	$.05 / (.20 + .05) = 0.2$
<del>clear</del>	<del>very cold</del> 0.25
snowing	mild 0.05
<del>snowing</del>	<del>cold</del> 0.10
<del>snowing</del>	<del>very cold</del> 0.10

# Chain Rule

**Definition: conditional probability**

$$P(h \mid e) = \frac{P(h, e)}{P(e)}$$

- We can run this **in reverse** to get

$$P(h, e) = P(h \mid e) \times P(e)$$

**Definition: chain rule**

$$\begin{aligned} P(\alpha_1, \dots, \alpha_n) &= P(\alpha_1) \times P(\alpha_2 \mid \alpha_1) \times \dots \times P(\alpha_n \mid \alpha_1, \dots, \alpha_{n-1}) \\ &= \prod_{i=1}^n P(\alpha_i \mid \alpha_1, \dots, \alpha_{i-1}) \end{aligned}$$


# Bayes' Rule

- From the **chain rule**, we have

$$\begin{aligned}P(h, e) &= P(h | e)P(e) \\ &= P(e | h)P(h)\end{aligned}$$

- Often**,  $P(e | h)$  is easier to compute than  $P(h | e)$ .

**Bayes' Rule:**



# Expected Value


- The **expected value** of a **function**  $f$  on a random variable is the weighted **average** of that function over the domain of the random variable, **weighted** by the **probability** of each value:

$$\mathbb{E} [f(X)] = \sum_{x \in \text{dom}(X)} P(X = x) f(x)$$


- The **conditional expected value** of a **function**  $f$  is the average value of the function over the domain, weighted by the **conditional probability** of each value:

$$\mathbb{E} [f(X) \mid Y = y] = \sum_{x \in \text{dom}(X)} P(X = x \mid Y = y) f(x)$$

# Expected Value Examples



$$\mathbb{E}[X] = 3$$
$$\mathbb{E}[X^2] \simeq 10$$



$$\mathbb{E}[X] = 3$$
$$\mathbb{E}[X^2] \simeq 12$$

# Summary

- **Probability** is a **numerical** measure of **uncertainty**
- Formal semantics:
  - Weights over **possible worlds** sum to 1
  - Probability of a proposition is **total weight** of **possible worlds** in which that proposition is **true**
- **Conditional probability** updates beliefs based on **evidence**
- **Expected value** of a function is its **probability-weighted average** over possible worlds