

Causality

CMPUT 366: Intelligent Systems

Bar §3.4

Lecture Outline


1. Recap
2. Causality Introduction
3. Causal Queries

Recap: Independence in a Belief Network


Belief Network Semantics:

Every node is **independent** of its **non-descendants**, **conditional only** on its **parents**


Patterns of dependence:



1. **Chain:** **Ends** are **not marginally** independent, but **conditionally** independent given middle



2. **Common ancestor:** **Descendants** are **not marginally** independent, but **conditionally** independent given ancestor



3. **Common descendant:** **Ancestors** are **marginally** independent, but **not conditionally** independent given descendant

Recap: Variable Elimination

1. Condition on observations by **conditioning**
2. Construct joint distribution factor by **multiplication**
3. Remove non-query, non-observed variables by **summing out**
4. **Normalize** at the end


Interleaving order of sums and products can improve **efficiency**:

$$\sum_A \sum_E f_1(Q, A, B, C) \times f_2(C, D, E) \quad \mathbf{112} \text{ computations}$$

$$= \left(\sum_A f_1(Q, A, B, C) \right) \times \left(\sum_E f_2(C, D, E) \right) \quad \mathbf{28} \text{ computations}$$

Causality Introduction: A Tale of Two Belief Networks

	Rain	Wet	P(Rain, Wet)
$P(\text{Rain} = T) = 0.5$	F	T	0.125
	F	F	0.375
$P(\text{Rain} = F) = 0.5$	T	T	0.45
	T	F	0.05




Natural network

- Two different ways to **factor** the joint distribution between whether the sidewalk is **Wet** and whether it is **Raining**:


$$\begin{aligned} P(\text{Rain}, \text{Wet}) &= P(\text{Wet} \mid \text{Rain})P(\text{Rain}) \\ &= P(\text{Rain} \mid \text{Wet})P(\text{Wet}) \end{aligned}$$

- Each factorization corresponds to a different **Belief Network**



Inverted network

The Inverted Network Isn't Crazy




Inverted network

Corresponds to the factoring $P(Rain | Wet)P(Wet)$

- Sometimes you want to answer the question
Given that I observe that the sidewalk is Wet, what is the probability that it is currently Raining?
- This is just updating our confidence in a hypothesis (it is **Raining**) given our observations (**Wet** sidewalk)
- Could **preprocess** the natural network into this form to avoid having to do a lot of computations with **Bayes' Rule**

The Inverted Network Is Crazy



Inverted network

Corresponds to the factoring $P(Rain | Wet)P(Wet)$

- If I **cause** my sidewalk to be **Wet** (by throwing water on it), what is the probability that it is **Raining**?

- So, condition on **Wet**=true
- This network seems to imply that it will be
 $P(Rain | Wet = True) = .78 > P(Rain) = .5$

- wait, what?

- **Question:** What is going wrong in this example?

Wet	P(Wet)
T	0.575
F	0.425

Rain	Wet	P(Rain Wet)
F	F	0.88
T	F	0.12
F	T	0.22
T	T	0.78

Observations vs. Interventions

- The semantics of Belief Networks are defined for **observational** questions
 - They don't directly model **causal** questions
 - In fact, in our Rainy Sidewalk example, we would get **exactly the same** (crazy) answer to our causal question from querying the **natural network**
- The joint distribution represented by the networks **doesn't model** the situation in which I **intervene**
 - Adding a variable **James_Throws_Water** to the distribution

Simpson's Paradox

Suppose we have information from two trials of a new drug:
One on male test subjects, and one on female test subjects.

G - gender
 D - received drug
 R - recovered

G	D	R	count	P(G,D,R)
M	T	T	18	0.225
M	T	F	12	0.15
M	F	T	7	0.0875
M	F	F	3	0.0375
F	T	T	2	0.025
F	T	F	8	0.1
F	F	T	9	0.1125
F	F	F	21	0.2625

- Is the drug **effective for males**?

$$P(R \mid D = \text{true}, G = \text{male}) = 0.60$$

$$P(R \mid D = \text{false}, G = \text{male}) = 0.70$$

- Is the drug **effective for females**?

$$P(R \mid D = \text{true}, G = \text{female}) = 0.20$$

$$P(R \mid D = \text{false}, G = \text{female}) = 0.30$$

- Is the drug **effective**?

$$P(R \mid D = \text{true}) = 0.50$$


$$P(R \mid D = \text{false}) = 0.40$$

Simpson's Paradox, explained

- The joint distribution factors as

$$P(G, D, R) = P(R | D, G) \times P(D | G) \times P(G)$$

- Per-gender** queries are answered **directly** by $P(R | D, G)$



- For the **overall query**, we want
$$P(R | D) = \frac{\sum_G P(R | G, D)P(G)}{\sum_{G,R} P(R | G, D)P(G)}$$

- But that's not how the distribution factors. If we follow the factoring above, we will instead compute



$$P(R | D) = \frac{P(R, D)}{P(D)} = \frac{\sum_G P(G, D, R)}{\sum_{G,R} P(G, D, R)} = \frac{\sum_G P(R | D, G)P(D | G)P(G)}{\sum_{G,R} P(R | D, G)P(D | G)P(G)}$$

- In our dataset, knowing whether a subject **got the drug** tells you something about their **gender**, and males have a **higher overall recovery** rate than females

- $P(R | G = \text{male}) = 0.625$ vs $P(R | G = \text{female}) = 0.275$

Selection Bias

- This problem is an example of **selection bias**
- Whether subjects received treatment is **systematically related** to their **response** to the treatment
- This is why **randomized trials** are the gold standard for causal questions:
 - The only thing that determines whether or not a subject is treated is a **random number**
 - Random number is definitely independent of **anything else** (including **response** to treatment)





Post-Intervention Distribution

- The causal query is really a query on a **different distribution** in which we have **forced** $D = true$
 - Different from the original joint distribution **conditioned** on **observing** that $D = true$
 - We will refer to the two distributions as the **observational** distribution and the **post-intervention** distribution
- With a post-intervention distribution, we can compute the answers to causal queries using **existing techniques** (e.g., variable elimination)

Post-Intervention Distribution for Simpson's Paradox

- **Observational distribution:**
$$P(G, D, R) = P(R \mid D, G) \times P(D \mid G) \times P(G)$$
- **Question:** What is the **post-intervention distribution** for Simpson's Paradox?
 - We're **forcing** $D = true$, so $P(D = true \mid G) = 1$ for all $g \in dom(G)$
 - That's the same as just omitting the $P(D \mid G)$ factor
- **Post-intervention distribution:**
$$P(G, D, R) = P(R \mid D, G) \times P(G)$$



The Do-Calculus

- How should we **express** causal queries?
- One approach: The **do-calculus**
- Condition on **observations**:
 $P(Y \mid X = x)$
- Express **interventions** with special **do** operator:
 $P(Y \mid do(X = x))$
- Allows us to **mix** observational and interventional information:
 $P(Y \mid Z = z, do(X = x))$

Evaluating Causal Queries With the Do-Calculus

Given a query $P(Y \mid do(X = x), Z = z)$:

1. Construct **post-intervention** distribution \hat{P} by **removing** all links from X 's direct parents to X
2. Evaluate the **observational** query $\hat{P}(Y \mid X = x, Z = z)$ in the **post-intervention distribution**

Example: Simpson's Paradox

- **Observational distribution:** $P(G,D,R) = P(R | D, G) \times P(D | G) \times P(G)$

- **Observational query:**

$$P(R | D) = \frac{P(R, D)}{P(D)} = \frac{\sum_G P(G, D, R)}{\sum_{G,R} P(G, D, R)} = \frac{\sum_G P(R | D, G) P(D | G) P(G)}{\sum_{G,R} P(R | D, G) P(D | G) P(G)}$$

- Observational query values: $P(R | D=true) = 0.50$ $P(R | D=false) = 0.40$

- **Post-intervention distribution** for causal query $P(R | do(D=true))$:



$$\hat{P}(G,D,R) = P(R | D, G) \times P(G)$$

- **Causal query:**

$$P(R | do(D = true)) = \hat{P}(R | D = true) = \frac{\sum_G P(R | D, G) P(G)}{\sum_{G,R} P(R | D, G) P(G)}$$

- Causal query values:

$$P(R | do(D=true)) = 0.40 \quad P(R | do(D=false)) = 0.50$$




Example: Rainy Sidewalk

Query: $P(\text{Rain} \mid \text{do}(\text{Wet}=\text{true}))$


Natural network:

- Observational distribution: $P(\text{Wet}, \text{Rain}) = P(\text{Wet} \mid \text{Rain})P(\text{Rain})$
- Post intervention distribution: $\hat{P}(\text{Wet}=\text{true}, \text{Rain}) = P(\text{Rain})P(\text{Wet})$
- $P(\text{Rain} \mid \text{do}(\text{Wet}=\text{true})) = .50$



Inverted network:

- Observational distribution: $P(\text{Wet}, \text{Rain}) = P(\text{Rain} \mid \text{Wet})P(\text{Rain})$
- Post intervention distribution:
 $\hat{P}(\text{Wet}=\text{true}, \text{Rain}) = P(\text{Rain} \mid \text{Wet})P(\text{Wet})$
- $P(\text{Rain} \mid \text{do}(\text{Wet}=\text{true})) = .78$



Causal Models

- The **natural network** gives the correct answer to our causal query, but the **inverted network** does not (**Why?**)
- Not every factoring of a joint distribution is a valid causal model

Definition:

A **causal model** is a directed acyclic graph of random variables such that for every edge $X \rightarrow Y$, the value of random variable X is **realized before** the value of random variable Y .



Alternative Representation: Influence Diagrams

Instead of adding a new operator, we can instead represent causal queries by **augmenting** the causal model with **decision variables** F_D for each potential intervention target D .

$$\text{dom}(F_D) = \text{dom}(D) \cup \{idle\}$$

$$P(D | pa(D), F_D) = \begin{cases} P(D | pa(D)) & \text{if } F_D = \textit{idle}, \\ 1 & \text{if } F_D \neq \textit{idle} \wedge D = F_D, \\ 0 & \text{otherwise.} \end{cases}$$

Influence Diagrams Examples



Summary

- **Observational** queries $P(Y \mid X=x)$ are different from **causal** queries $P(Y \mid \text{do}(X=x))$
- To evaluate **causal query** $P(Y \mid \text{do}(X=x))$:
 1. Construct **post-intervention distribution** \hat{P} by **removing** all links from X 's direct parents to X
 2. Evaluate the **observational** query $\hat{P}(Y \mid X=x, Z=z)$ in the **post-intervention distribution**
- Not every correct Bayesian network is a valid causal model