

Conditional Independence

CMPUT 366: Intelligent Systems

P&M §8.2

Logistics & Assignment #1

- **Assignment #1** is due Feb 4 11:59pm (**next week**)
- Office hours have begun!
 - Not mandatory; for getting help from TAs
 - There are no labs for this course

Lecture Outline

1. Recap
2. Expected Value
3. Structure
4. Marginal Independence
5. Conditional Independence

Recap: Probability

- **Probability** is a numerical measure of **uncertainty**
 - **Not** a measure of **truth**
- **Semantics:**
 - A **possible world** is a **complete assignment** of values to variables
 - Every possible world has a probability
 - Probability of a **proposition** is the sum of probabilities of **possible worlds** in which the statement is **true**

Recap:

Conditional Probability

- When we receive **evidence** in the form of a proposition e , it **rules out** all of the possible worlds in which e is **false**
 - We set those worlds' probability to **0**, and **rescale** remaining probabilities to sum to **1**
- Result is probabilities **conditional on e** : $P(h \mid e)$

Expected Value


- The **expected value** of a **function** f on a random variable is the weighted **average** of that function over the domain of the random variable, **weighted** by the **probability** of each value:

$$\mathbb{E} [f(X)] = \sum_{x \in \text{dom}(X)} P(X = x) f(x)$$


- The **conditional expected value** of a **function** f is the average value of the function over the domain, weighted by the **conditional probability** of each value:

$$\mathbb{E} [f(X) \mid Y = y] = \sum_{x \in \text{dom}(X)} P(X = x \mid Y = y) f(x)$$

Expected Value Examples



$$\mathbb{E}[X] = 3$$
$$\mathbb{E}[X^2] \simeq 10$$



$$\mathbb{E}[Y] = 3$$
$$\mathbb{E}[Y^2] \simeq 12$$

Unstructured Joint Distributions

- Probabilities are not fully **arbitrary**
 - **Semantics** tell us probabilities given the joint distribution.
 - Semantics alone do not restrict probabilities **very much**
- In general, we might need to **explicitly** specify the entire **joint distribution**
 - **Question:** How many numbers do we need to assign to fully specify a joint distribution of k binary random variables?
- We call situations where we have to explicitly enumerate the entire joint distribution **unstructured**

Structure

- Unstructured domains are very hard to reason about
- Fortunately, most real problems are generated by some sort of **underlying process**
 - This gives us **structure** that we can exploit to represent and reason about probabilities in a more **compact** way
 - We can **compute** any required joint probabilities based on the process, instead of specifying every possible joint probability explicitly
- Simplest kind of structure is when random variables don't **interact**

Marginal Independence

When the value of one variable **never** gives you information about the value of the other, we say the two variables are **marginally independent**.

Definition:

Random variables X and Y are **marginally independent** iff

1. $P(X = x \mid Y = y) = P(X = x)$, and
2. $P(Y = y \mid X = x) = P(Y = y)$

for all values of $x \in \text{dom}(X)$ and $y \in \text{dom}(Y)$.

Marginal Independence Example

- I flip four **fair coins**, and get four results: C_1, C_2, C_3, C_4
- **Question:** What is the probability that C_1 is **heads**?
 - $P(C_1 = heads)$
- Suppose that $C_2, C_3,$ and C_4 are **tails**
- **Question:** Now what is the probability that C_1 is **heads**?
 - $P(C_1 = heads \mid C_2 = tails, C_3 = tails, C_4 = tails)$
 - Why?

Properties of Marginal Independence

Proposition:

If X and Y are marginally independent, then

$$P(X = x, Y = y) = P(X = x)P(Y = y)$$

for all values of $x \in \text{dom}(X)$ and $y \in \text{dom}(Y)$.

Proof:

1. $P(X = x, Y = y) = P(X = x | Y = y)P(Y = y)$ Chain rule

2. $P(X = x, Y = y) = P(X = x)P(Y = y)$ Marginal independence



Exploiting Marginal Independence

C ₁	P
H	0.5

C ₂	P
H	0.5

C ₃	P
H	0.5

C ₄	P
H	0.5

- Instead of storing the **entire joint distribution**, we can store 4 **marginal distributions**, and use them to recover joint probabilities
 - **Question:** How many numbers do we need to assign to fully specify the marginal distribution for a **single** binary variable?
- If **everything** is independent, learning from observations is hopeless (**why?**)
 - But also if **nothing** is independent
 - The **intermediate** case, where many variables are independent, is ideal

C ₁	C ₂	C ₃	C ₄	P
H	H	H	H	0.0625
H	H	H	T	0.0625
H	H	T	H	0.0625
H	H	T	T	0.0625
H	T	H	H	0.0625
H	T	H	T	0.0625
H	T	T	H	0.0625
H	T	T	T	0.0625
T	H	H	H	0.0625
T	H	H	T	0.0625
T	H	T	H	0.0625
T	H	T	T	0.0625
T	T	H	H	0.0625
T	T	H	T	0.0625
T	T	T	H	0.0625

Clock Scenario


Example:

- I have a stylish but impractical clock with no number markings
- Two students, Alice and Bob, both look at the clock at the same time, and form opinions about what time it is
 - Their opinion of the time is **directly affected** by the actual time
 - They don't talk to each other, so Alice's opinion of the time is not directly affected by Bob's opinion of the time (& vice versa)
- **Question:** Are A and B **marginally independent**?

$$P(A | B) \neq P(A)$$

- **Question:** If we know it is 10:09. Are A and B **independent**?

$$P(A | B, T = 10:09) = P(A | T = 10:09)$$



Random variables:

A - Time Alice thinks it is

B - Time Bob thinks it is

T - Actual time

Conditional Independence

When knowing the value of a **third** variable Z **makes** two variables A, B independent, we say that they are **conditionally independent given Z** (or **independent conditional on Z**).

Definition:

Random variables X and Y are **conditionally independent given Z** iff

$$P(X = x \mid Y = y, Z = z) = P(X = x \mid Z = z)$$

for all values of $x \in \text{dom}(X)$, $y \in \text{dom}(Y)$, and $z \in \text{dom}(Z)$.

We write this using the notation $X \perp\!\!\!\perp Y \mid Z$.

Clock example: A and B are conditionally independent given T .

Properties of Conditional Independence

Proposition:

If X and Y are conditionally independent given Z , then

$$P(X = x, Y = y \mid Z) = P(X = x \mid Z)P(Y = y \mid Z)$$

for all values of $x \in \text{dom}(X)$, $y \in \text{dom}(Y)$, and $z \in \text{dom}(Z)$.

Proof:

1. $P(X = x, Y = y \mid Z) = P(X = x \mid Y = y, Z = z)P(Y = y \mid Z)$ Chain rule

2. $P(X = x, Y = y \mid Z) = P(X = x \mid Z)P(Y = y \mid Z)$ Conditional independence



Properties of Conditional Independence

Question: Is conditional independence **commutative**?

- i.e., If $X \perp\!\!\!\perp Y \mid Z$, is it also true that $Y \perp\!\!\!\perp X \mid Z$?

Proof:

$$X \perp\!\!\!\perp Y \mid Z \iff P(X, Y \mid Z) = P(X \mid Z)P(Y \mid Z) \text{ previous result}$$

$$\iff P(Y, X \mid Z) = P(Y \mid Z)P(X \mid Z) \text{ commutativity of multiplication}$$

$$\iff Y \perp\!\!\!\perp X \mid Z \quad \blacksquare$$

Exploiting Conditional Independence

If X and Y are marginally independent given Z , then we can again just store **smaller tables** and recover joint distributions by **multiplication**.

- **Question:** How many **tables** do we need to store in order to be able to compute the joint distribution of X, Y, Z when X and Y are independent given Z ?
- i.e., how many table to be able to compute $P(X = x, Y = y, Z = z)$ for every combination of x, y, z ?

Preview: In the upcoming lectures, we will see how to efficiently exploit **complex structures** of conditional independence

Simplified Clock Example

A	T	P(A T)
12	1	0.25
1	1	0.50
2	1	0.25
1	2	0.25
2	2	0.50
3	2	0.25
2	3	0.25
3	3	0.50
4	3	0.25
	⋮	
	⋮	
	⋮	

B	T	P(B T)
12	1	0.25
1	1	0.5
2	1	0.25
1	2	0.25
2	2	0.5
3	2	0.25
2	3	0.25
3	3	0.5
4	3	0.25
	⋮	
	⋮	
	⋮	

T	P(T)
1	0
2	1/10
3	1/10
4	1/10
5	1/10
6	1/10
7	1/10
8	1/10
9	1/10
10	1/10
11	1/10
12	0

$$\begin{aligned}
 &P(A = 1, B = 2, T = 2) \\
 &= P(A = 1 | T = 2)P(B = 2 | T = 2)P(T = 2) \\
 &= 0.25 \times 0.5 \times 0.10 \\
 &= 0.0125
 \end{aligned}$$

$$\begin{aligned}
 &P(A = 1, B = 2, T = 1) \\
 &= P(A = 1 | T = 1)P(B = 2 | T = 1)P(T = 1) \\
 &= 0.5 \times 0.25 \times 0.0 \\
 &= 0
 \end{aligned}$$

Caveats

- Often, when two variables are **marginally** independent, they are also **conditionally** independent given a third variable
 - E.g., coins C_1 , and C_2 are marginally independent, **and also** conditionally independent given C_3 : Learning the value of C_3 does not make C_2 any more informative about C_1 .
- This is **not always true**
 - Consider another random variable: B is true if both C_1 and C_2 are the **same** value
 - C_1 and C_2 are **marginally independent**: $P(C_1 = heads \mid C_2 = heads) = P(C_1 = heads)$
 - In fact, C_1 and C_2 are also both **marginally independent of B** : $P(C_1 \mid B = true) = P(C_1)$
 - But if I know the value of B , then knowing the value of C_1 tells me **exactly** what the value of C_2 must be: $P(C_1 = heads \mid B = true, C_2 = heads) \neq P(C_1 = heads \mid B = true)$
 - C_1 and C_2 are **not conditionally independent given B**

Summary

- **Unstructured** joint distributions are **exponentially** expensive to represent (and operate on)
- **Marginal and conditional independence** are especially important forms of **structure** that a distribution can have
 - Vastly **reduces the cost** of representation and computation
 - **Caveat:** The **relationship** between marginal and conditional independence is not fixed
- Joint probabilities of (conditionally or marginally) **independent** random variables can be computed by **multiplication**