

Probability Theory

CMPUT 296: Basics of Machine Learning

§2.1-2.2

Recap

This class is about **understanding** machine learning techniques by understanding their basic **mathematical underpinnings**

- Course details at jrwright.info/mlbasics/ and on eClass: <https://eclass.srv.ualberta.ca/course/view.php?id=64044>
- Exams will be **spot checked** but not proctored
- Readings in free textbook, with associated thought questions

Logistics

- Videos for Tuesday's and today's lectures will be released today on eClass
- Assignment 1 will be released today on eClass
- Thought Question 1 will be released today on eClass
- No TA office hours this week

Outline

1. Recap & Logistics
2. Probabilities
3. Defining Distributions
4. Random Variables

Why Probabilities?

Even if the world is completely deterministic, outcomes can **look random** (**why?**)

Example: A high-tech gumball machine behaves according to

$f(x_1, x_2) = \text{output candy if } x_1 \text{ \& } x_2,$

where $x_1 = \text{has candy}$ and $x_2 = \text{battery charged}$.

- You can only see if it has candy
- From your perspective, when $x_1 = 1$, sometimes candy is output, sometimes it isn't
- It **looks stochastic**, because it depends on the hidden input x_2

Measuring Uncertainty

- **Probability** is a way of **measuring** uncertainty
- We assign a number between 0 and 1 to **events** (hypotheses):
 - **0** means absolutely certain that statement is **false**
 - **1** means absolutely certain that statement is **true**
 - **Intermediate** values mean more or less certain
- Probability is a measurement of **uncertainty**, **not truth**
 - A statement with probability .75 is not "mostly true"
 - Rather, we **believe** it is more **likely** to be true than not

Subjective vs. Objective: The Frequentist Perspective

- Probabilities can be interpreted as **objective** statements about the **world**, or as **subjective** statements about an agent's **beliefs**.
- Objective view is called **frequentist**:
 - The probability of an event is the proportion of times it would happen **in the long run** of **repeated experiments**
 - Every event has a single, **true** probability
 - Events that can only happen **once** don't have a well-defined probability

Subjective vs. Objective: The Bayesian Perspective

- Probabilities can be interpreted as **objective** statements about the **world**, or as **subjective** statements about an agent's **beliefs**.
- Subjective view is called **Bayesian**:
 - The probability of an event is a measure of an agent's **belief** about its likelihood
 - Different agents can legitimately have **different beliefs**, so they can legitimately assign **different probabilities** to the same event
 - There is **only one way** to **update** those beliefs in response to new data

Prerequisites Check

- Derivatives
 - Rarely integration
 - I will teach you about partial derivatives
- Vectors, dot-products, matrices
- Set notation
 - Complement A^c of a set, union $A \cup B$ of sets, intersection of sets $A \cap B$
 - Set of sets, power set $\mathcal{P}(A)$
- Basics of probability. (We will refresh today)

Terminology

- If you are unsure, notation sheet in the notes is a good starting point
- **Countable:** A set whose elements can be assigned an integer index
 - The integers themselves
 - Any finite set, e.g., $\{0.1, 2.0, 3.7, 4.123\}$
 - We'll sometimes say **discrete**, even though that's a little imprecise
- **Uncountable:** Sets whose elements *cannot* be assigned an integer index
 - Real numbers \mathbb{R}
 - Intervals of real numbers, e.g., $[0, 1]$, $(-\infty, 0)$
 - Sometimes we'll say **continuous**

Outcomes and Events

All probabilities are defined with respect to a **measurable space** (Ω, \mathcal{E}) of **outcomes** and **events**:

- Ω is the **sample space**: The set of all possible outcomes
- $\mathcal{E} \subseteq \mathcal{P}(\Omega)$ is the **event space**: A set of subsets of Ω satisfying

1. $A \in \mathcal{E} \implies A^c \in \mathcal{E}$

2. $A_1, A_2, \dots \in \mathcal{E} \implies \bigcup_{i=1}^{\infty} A_i \in \mathcal{E}$

Event Spaces

Definition:

A set $\mathcal{E} \subseteq \mathcal{P}(\Omega)$ is an **event space** if it satisfies

$$1. A \in \mathcal{E} \implies A^c \in \mathcal{E}$$

$$2. A_1, A_2, \dots \in \mathcal{E} \implies \bigcup_{i=1}^{\infty} A_i \in \mathcal{E}$$

1. A collection of outcomes (e.g., either a 2 or a 6 were rolled) is an event.
2. If we can measure that an event has occurred, then we should also be able to measure that the event has not occurred; i.e., its **complement** is measurable.
3. If we can measure two events separately, then we should be able to tell if one of them has happened; i.e., their **union** should be measurable too.

Discrete vs. Continuous Sample Spaces

Discrete (countable) outcomes

$$\Omega = \{1,2,3,4,5,6\}$$

$$\Omega = \{\text{person, woman, man, camera, TV, ...}\}$$

$$\Omega = \mathbb{N}$$

$$\mathcal{E} = \{\emptyset, \{1,2\}, \{3,4,5,6\}, \{1,2,3,4,5,6\}\}$$

Typically: $\mathcal{E} = \mathcal{P}(\Omega)$

Question:

$$\mathcal{E} = \{\{1\}, \{2\}, \{3\}, \{4\}, \{5\}, \{6\}\}?$$

Continuous (uncountable) outcomes

$$\Omega = [0,1]$$

$$\Omega = \mathbb{R}$$

$$\Omega = \mathbb{R}^k$$

$$\mathcal{E} = \{\emptyset, [0,0.5], (0.5,1.0], [0,1]\}$$

Typically: $\mathcal{E} = B(\Omega)$ ("Borel field")

Note: *not* $\mathcal{P}(\Omega)$

Axioms

Definition:

Given a measurable space (Ω, \mathcal{E}) , any function $P : \mathcal{E} \rightarrow [0,1]$ satisfying

1. **unit measure:** $P(\Omega) = 1$, and

2. **σ -additivity:** $P\left(\bigcup_{i=1}^{\infty} A_i\right) = \sum_{i=1}^{\infty} P(A_i)$ for any countable sequence

$A_1, A_2, \dots \in \mathcal{E}$ where $A_i \cap A_j = \emptyset$ whenever $i \neq j$

is a **probability measure** (or **probability distribution**).

If P is a probability measure over (Ω, \mathcal{E}) , then (Ω, \mathcal{E}, P) is a **probability space**.

Defining a Distribution

Example:

$$\Omega = \{0,1\}$$

$$\mathcal{E} = \{\emptyset, \{0\}, \{1\}, \Omega\}$$

$$P = \begin{cases} 1 - \alpha & \text{if } A = \{0\} \\ \alpha & \text{if } A = \{1\} \\ 0 & \text{if } A = \emptyset \\ 1 & \text{if } A = \Omega \end{cases}$$

where $\alpha \in [0,1]$.

Questions:

1. Do you recognize this distribution?
2. How should we choose P in practice?
 - a. Can we choose an arbitrary function?
 - b. How can we guarantee that all of the constraints will be satisfied?

Probability Mass Functions (PMFs)

Definition: Given a **discrete** sample space Ω and event space $\mathcal{E} = \mathcal{P}(\Omega)$, any function $p : \Omega \rightarrow [0,1]$ satisfying $\sum_{\omega \in \Omega} p(\omega) = 1$ is a **probability mass function**.

- For a discrete sample space, instead of defining P directly, we can define a **probability mass function** $p : \Omega \rightarrow [0,1]$.
- p gives a probability for **outcomes** instead of **events**
- The probability for any event $A \in \mathcal{E}$ is then defined as $P(A) = \sum_{\omega \in \Omega} p(\omega)$.

Example: PMF for a Fair Die

A **categorical distribution** is a distribution over a **finite** outcome space, where the probability of each outcome is specified separately.

Example: Fair Die

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

$$p(\omega) = \frac{1}{6}$$


ω	$p(\omega)$
1	1/6
2	1/6
3	1/6
4	1/6
5	1/6
6	1/6

Questions:

1. What is a possible event?
What is its probability?
2. What is the event space?

Example: Using a PMF

- Suppose that you recorded your commute time (in minutes) every day for a year (i.e., 365 recorded times).
- **Question:** How do you get $p(t)$?
- **Question:** How is $p(t)$ useful?



Useful PMFs: Bernoulli

A **Bernoulli distribution** is a special case of a **categorical distribution** in which there are only two outcomes. It has a single **parameter** $\alpha \in (0,1)$.

$$\Omega = \{T, F\} \text{ (or } \Omega = \{S, F\})$$

$$\text{Alternatively: } \Omega = \{0,1\}$$


$$p(\omega) = \begin{cases} \alpha & \text{if } \omega = T \\ 1 - \alpha & \text{if } \omega = F. \end{cases}$$

$$p(k) = \alpha^k(1 - \alpha)^{1-k} \text{ for } k \in \{0,1\}$$

Useful PMFs: Poisson

A **Poisson distribution** is a distribution over the non-negative integers. It has a single parameter $\lambda \in (0, \infty)$.

E.g., number of calls received by a call centre in an hour, number of letters received per day.



$$p(k) = \frac{\lambda^k e^{-\lambda}}{k!}$$


Questions:

1. Could we define this with a table instead of an equation?
2. How can we check whether this is a valid PMF?


Commute Times Again

- **Question:** Could we use a **Poisson distribution** for commute times (instead of a categorical distribution)?
- **Question:** What would be the benefit of using a Poisson distribution?

$$p(k) = \frac{\lambda^k e^{-\lambda}}{k!}$$




$$p(4) = 1/365, p(5) = 2/365, p(6) = 4/365, \dots$$



Continuous Commute Times

- It never actually takes *exactly* 12 minutes; I rounded each observation to the nearest integer number of minutes.
- Actual data was 12.345 minutes, 11.78213 minutes, etc.
- **Question:** Could we use a Poisson distribution to predict the *exact* commute time (rather than the nearest number of minutes)? Why?



Probability Density Functions (PDFs)

Definition: Given a **continuous** sample space Ω and event space $\mathcal{E} = B(\Omega)$, any function $p : \Omega \rightarrow [0, \infty)$ satisfying $\int_{\Omega} p(\omega) d\omega = 1$ is a **probability density function**.

- For a continuous sample space, instead of defining P directly, we can define a **probability density function** $p : \Omega \rightarrow [0, \infty)$.
- The probability for any event $A \in \mathcal{E}$ is then defined as

$$P(A) = \int_A p(\omega) d\omega.$$

PMFs vs PDFs

1. When sample space Ω is **discrete**:

- Singleton event: $P(\{\omega\}) = p(\omega)$ for $\omega \in \Omega$

$$P(A) = \sum_{\omega \in \Omega} p(\omega)$$

2. When sample space Ω is **continuous**:

- Example: Stopping time for a car with $\Omega = [3, 12]$
- **Question:** What is the probability that the stopping time is *exactly* 3.14159?

$$P(A) = \int_A p(\omega) d\omega$$

$$P(\{3.14159\}) = \int_{3.14159}^{3.14159} p(\omega) d\omega$$


- More reasonable: Probability that stopping time is between 3 to 3.5.

Useful PDFs: Uniform

A **uniform distribution** is a distribution over a real interval. It has two parameters: a and b .

$$\Omega = [a, b]$$

$$p(\omega) = \begin{cases} \frac{1}{b-a} & \text{if } a \leq \omega \leq b, \\ 0 & \text{otherwise.} \end{cases}$$



Question: Does Ω have to be bounded?


Useful PDFs: Gaussian

A **Gaussian distribution** is a distribution over the real numbers. It has two parameters: $\mu \in \mathbb{R}$ and $\sigma \in \mathbb{R}^+$.

$$\Omega = \mathbb{R}$$

$$p(\omega) = \frac{1}{\sqrt{2\pi\sigma^2}} \exp\left(-\frac{1}{2\sigma^2}(\omega - \mu)^2\right)$$

where $\exp(x) = e^x$




Useful PDFs: Exponential

An **exponential distribution** is a distribution over the positive reals. It has one parameter $\lambda > 0$.

$$\Omega = \mathbb{R}$$

$$p(\omega) = \lambda \exp(-\lambda\omega)$$



Why can the density be above 1?

Consider an interval event $A = [x, x + \Delta x]$, for small Δx .

$$P(A) = \int_x^{x+\Delta x} p(\omega) d\omega$$
$$\approx p(x)\Delta x$$

- $p(x)$ can be big, because Δx can be very small
 - In particular, $p(x)$ can be bigger than 1
- But $P(A)$ **must** be less than or equal to 1

Random Variables

Random variables are a way of reasoning about a complicated underlying probability space in a more straightforward way.

Example: Suppose we observe both a die's number, and where it lands.

$$\Omega = \{(left,1), (right,1), (left,2), (right,2), \dots, (right,6)\}$$

We might want to think about the probability that we get a large number, without thinking about where it landed.

We could ask about $P(X \geq 4)$, where

X = number that comes up.

Random Variables, Formally

Given a probability space (Ω, \mathcal{E}, P) , a **random variable** is a function $X : \Omega \rightarrow \Omega_X$ (where Ω_X is some other outcome space), satisfying

$$\{\omega \in \Omega \mid X(\omega) \in A\} \in \mathcal{E} \quad \forall A \in B(\Omega_X).$$

It follows that $P_X(A) = P(\{\omega \in \Omega \mid X(\omega) \in A\})$.

Example: Let Ω be a population of people, and $X(\omega) = \text{height}$, and $A = [5'1'', 5'2'']$.

$$P(X \in A) = P(5'1'' \leq X \leq 5'2'') = P(\{\omega \in \Omega : X(\omega) \in A\}).$$

Random Variables and Events

- A Boolean expression involving random variables defines an event:

$$\text{E.g., } P(X \geq 4) = P(\{\omega \in \Omega \mid X(\omega) \geq 4\})$$


- Similarly, every event can be understood as a Boolean random variable:

$$Y = \begin{cases} 1 & \text{if event } A \text{ occurred} \\ 0 & \text{otherwise.} \end{cases}$$

- From this point onwards, we will exclusively reason in terms of random variables rather than probability spaces.

Example: Histograms

Consider the continuous commuting example again, with observations 12.345 minutes, 11.78213 minutes, etc.



- **Question:** What is the random variable?
- **Question:** How could we turn our observations into a histogram?

Summary

- Probabilities are a means of **quantifying uncertainty**
- A probability distribution is defined on a measurable space consisting of a **sample space** and an **event space**.
- **Discrete** sample spaces (and random variables) are defined in terms of **probability mass functions** (PMFs)
- **Continuous** sample spaces (and random variables) are defined in terms of **probability density functions** (PDFs)
- **Random variables** are more convenient than operating directly on probability spaces